CELL-DYN 26 Plus Control








IMPORTANT: The barcode is for use only on the CELL-DYN Ruby. Refer to the appropriate System Operator's Manual for proper use of CELL-DYN Calibrator and Control Products. IMPORTANT: Mixing and Handling

- 1. Remove a vial of the control from the refrigerator and warm to room temperature (18° to 30° C) for 15 minutes before use.
- 2. To mix: (Do NOT mix mechanically or vortex.)

For a video demonstration, visit www.corelaboratory.abbott and navigate to the Customer Portal → Technical Library → Other Reference Documents → Hematology Aids.

a. Hold the vial vertically and roll each vial between the palms of the hands for 15-20 seconds.



b. Continue to mix by holding the vial by the ends between the thumb and finger, rapidly inverting the vial 20 times end-over-end using a very quick turning motion of the wrist.



- c. Analyze immediately after mixing. Subsequent analyses during this test period may be performed by inverting the vial 5 times prior to instrument analysis.
- d. Steps a-c must be repeated upon removing the sample from the refrigerator for the entire open-vial time period regardless of the method of analysis (open tube, cap piercing, auto sample or manual sample).
- 3. Refer to the appropriate CELL-DYN System Operator's Manual for information about analyzing control specimens.
- 4. FOR AUTOMATED SAMPLING OR MANUAL CLOSED SAMPLING (CS):
 - Refer to the appropriate CELL-DYN Operator's Manual. Remove the vial from the sample handler immediately after sampling.

FOR OPEN-VIAL SAMPLING:

- Aspirate a sample from the vial.
- Carefully wipe the vial rim and cap with a lint-free tissue.
- Replace the cap, ensuring it is on tight.

After sampling, return vial to refrigerator for maximum open-vial stability. If run in the open mode, wipe the threads of both vial and cap before replacing cap and returning to refrigerator.

CELL-DYN 26 Plus Control



Exp. 2024-04-05 8 Consecutive Day Open-Vial Stability CONTROL H CONTROL L CONTROL N LOT L4022 LOT N4022 LOT H4022 SYSTEM **ASSAY VALUE** ± MEAN RANGE * **ASSAY VALUE** ± MEAN RANGE * **ASSAY VALUE** ± MEAN RANGE * PARAMETER N/A N/A N/A N/A WOC 109/L N/A WIC 109/L N/A N/A N/A N/A N/A N/A WBC 109/L N/A N/A N/A N/A N/A N/A NEU 109/L NEU % N/A N/A N/A N/A N/A N/A LYM 109/L N/A N/A N/A N/A N/A N/A LYM % N/A N/A N/A N/A N/A N/A MONO 10º/L N/A N/A N/A N/A N/A N/A EOS 109/L N/A N/A N/A N/A N/A N/A CELL-DYN 3700 EOS % N/A N/A N/A N/A N/A N/A SYSTEM BASO 109/L N/A N/A N/A N/A N/A N/A BASO % N/A N/A N/A N/A N/A N/A HGB g/dL N/A N/A N/A N/A N/A N/A HCT % N/A N/A N/A N/A N/A N/A MCV fL N/A N/A N/A N/A N/A N/A MCH pg N/A N/A N/A N/A N/A N/A MCHC g/dL RDW % N/A N/A N/A N/A N/A N/A PLT 10°/L N/A N/A N/A N/A N/A N/A MPV fL N/A N/A N/A N/A N/A N/A WBC (WOC) 10°/L 3.9 0.4 6.6 0.7 15.2 2.5 WBC (NOC) 109/L 4.1 0.4 6.9 1.0 15.7 2.5 0.3 0.8 8.5 NEU 109/L 2.2 3.6 2.0 NEU % 55.0 8.0 54.9 8.0 55.8 10.0 LYM 109/L 1.1 0.3 2.0 0.8 4.4 2.0 LYM % 28.8 9.0 29.7 9.0 10.0 29.1 MONO 109/L 0.4 0.2 0.6 0.4 1.4 0.6 10.0 5.0 5.0 9.4 MONO % 9.7 3.0 EOS 109/L 0.1 0.1 0.2 0.2 0.4 0.2 **CELL-DYN Ruby** 3.0 3.0 2.5 2.0 2.6 1.0 BASO 109/L 0.1 0.1 0.2 0.6 0.2 0.6 SYSTEM 3.0 3.6 3.0 3.1 3.1 3.0 BASO % RBC 1012/L 2.99 0.15 4.13 0.20 5.30 0.30 HGB g/dL 7.4 0.4 11.2 0.6 15.6 8.0 HCT % 20.7 1.5 30.8 2.5 41.3 3.5 MCV fL 69.3 4.0 74.5 4.0 77.9 5.0 27.1 24.6 2.0 2.0 29.5 2.0 MCH pg 2.3 3.0 37.9 MCHC g/dL 35.5 36.4 2.3 2.5 2.5 2.5 RDW % 14.6 12.6 11.2 PLT 109/L 76 20 225 30 545 60 10.1 3.0 9.9 3.0 9.9 3.0




CELL-DYN and CELL-DYN Ruby are trademarks of Abbott Laboratories in various jurisdictions.



Abbott Laboratories Diagnostics Division Abbott Park, IL 60064 EC REP Abbott GmbH & Co. KG Max-Planck-Ring 2

MANUFACTURED FOR Abbott Laboratories 65205 Wiesbaden Germany +49-6122-580









The **MEAN RANGE** does not represent standard deviations (SD). NOTE: Flags may occur with control materials and should be disregarded.